


4-1BBL mRNA-LNP

Ready-to-use lipid nanoparticles



Products and Services

- Monoclonal Antibody
- Bispecific Antibody
- mRNA-LNP
- Hybridoma Sequencing
- CAR-T/NK
- Stable Cell Line

Order Information

Catalog#	PM-LNP-0143	Size	200uL
----------	-------------	------	-------

Description

4-1BBL (4-1BB ligand), also known as tumor necrosis factor ligand superfamily member 9 (TNFSF9), is a type II membrane protein expressed by T cells. Binding of 4-1BBL to its receptor 4-1BB (also known as CD137) provides co-stimulatory signals to the T cells, promoting cell proliferation and survival. 4-1BBL can also enhance TCR/CD3-triggered activation-induced T cell apoptosis, modulate CD28 co-stimulation to promote Th1 cell responses, and induce peripheral monocyte proliferation. 4-1BBL is comprised of 254 amino acids and its GenPept accession number is NP_003802. ProMab's PM-LNP-0060 nanoparticles contain the 4-1BBL mRNA protected by a lipid shell. The nanoparticles are formulated with SM-102, DSPC, cholesterol and DMG-PEG2000 at an optimal molar concentration for a high rate of encapsulation and efficient mRNA delivery in vitro and in vivo.

Composition

mRNA-LNPs suspended in PBS (-Ca, -Mg) (pH: 7.0-7.4).

Storage

Product is delivered on wet ice. Store at 4°C for up to 3 months.

Handling

Upon receipt, centrifuge the vial for a few seconds to ensure the contents are located at the bottom of the vial. Vortex mixing or prolonged centrifugation may rupture the nanoparticles. Store the vial of nanoparticles in the refrigerator and keep on ice when in use. Do not allow the nanoparticles to warm to room temperature. mRNA-LNP suspensions should only be handled with certified RNase-free reagents and consumables. The use of filtered pipette tips is highly recommended.

4-1BBL mRNA-LNP

Ready-to-use lipid nanoparticles

Safety & Research Disclosure

All ProMab mRNA lipid nanoparticle products are for in vitro research use only. Products are not FDA approved for human use.

Protocol for Transfecting Suspension Cells

Suspend 0.5 - 1 million cells in 1 ml of culture medium. Ensure the cells are healthy and well-dispersed, as cell clumping may reduce transfection efficiency. Disperse the nanoparticle suspension by gently pipetting up and down several times, then slowly add 20-40 μ l to the cells, dropwise. Gently mix the cells and incubate them overnight in a culture incubator. The next day, and every day thereafter, check the culture for expression of the protein encoded by the mRNA-LNP. Cell-bound proteins can be detected by flow cytometry or western blotting using the transfected cells, whereas secreted proteins can be detected by ELISA, western blotting or flow cytometry (on a target cell line) using the medium collected from the transfected cells.

4-1BBL Expression in HEK293S Cells Treated with PM-LNP-0143




Figure 1. Flow Cytometry. PM-LNP-0143 nanoparticle-treated HEK293S cells express 4-1BBL, detected with an anti-4-1BBL antibody.

Products and Services

- Monoclonal Antibody
- Bispecific Antibody
- mRNA-LNP
- Hybridoma Sequencing
- CAR-T/NK
- Stable Cell Line