

A1847-GFP-LUC cells were generated from the human ovarian cancer cell line A1847 by transduction with replication-defective lentivirus encoding eGFP and luciferase. Expression of eGFP was confirmed by flow cytometry (Figure 1) and expression of luciferase was confirmed by luminescence after exposure to luciferin (Figure 2).

Storage
Store vial in liquid nitrogen immediately upon receipt.
Formulation
Cells are cryopreserved in 0.5 ml - 1 ml of 90% FBS + 10% DMSO.
Thaw Protocol
Partially immerse the vial in a 37°C water bath with gentle shaking until most of the medium is thawed. In a tissue culture hood, add 1 ml of pre-warmed culture medium into the vial and immediately transfer the contents of the vial to a centrifuge tube containing 5-10 ml of pre-warmed culture medium. Centrifuge the tube at room temperature for 5 minutes, aspirate the supernatant and suspend the cell pellet in 5-10 ml of pre-warmed culture medium.
Culture Protocol
A1847-GFP-LUC is an adherent cell line. Culture the cells in DMEM containing 10% FBS using a humidified incubator set to 5% CO ₂ . When the cell monolayer is nearly confluent, use trypsin-EDTA to detach the cells from the culture flask and immediately neutralize the trypsin by adding fresh, pre-warmed culture medium to the cells. Cells should be passaged at least twice per week.
Notices & Disclaimer
ProMab Biotechnologies products are intended for laboratory research purposes only, not for use in humans. This product is not for resale and may not be transferred to a third party without written consent from ProMab Biotechnologies, Inc. To purchase this product, you must accept the terms and conditions of ProMab Biotechnologies' Material Transfer Agreement (MTA).

All products are for research use only

Discover more | ProMab.com



📍 2600 Hilltop Dr, Building B, Richmond, CA 94806
☎ 1.866.339.0871 | ✉ info@promab.com
☎ 510.740.3625 | ✉ customerservice@promab.com






Figure 1. PM-A1847-GFP-LUC cells and parental A1847 cells were analyzed for GFP fluorescence by flow cytometry. Only the PM-A1847-GFP-LUC cells were fluorescent.




Figure 2. PM-A1847-GFP-LUC cells and parental A1847 cells were lysed and incubated with luciferin to assess luciferase expression. Only PM-A1847-GFP-LUC cells became luminescent.

All products are for research use only

Discover more | Promab.com



2600 Hilltop Dr, Building B, Richmond, CA 94806
1.866.339.0871 | info@promab.com
510.740.3625 | customerservice@promab.com

