

RPMI8226-GFP-LUC cells were generated from the human myeloma cell line RPMI8226 by transduction with replication-defective lentivirus encoding eGFP and luciferase. Expression of eGFP was confirmed by flow cytometry (Figure 1) and expression of luciferase was confirmed by luminescence after exposure to luciferin (Figure 2).

| |
|--|
| Storage |
| Store vial in liquid nitrogen immediately upon receipt. |
| Formulation |
| Cells are cryopreserved in 0.5 ml - 1 ml of 90% FBS + 10% DMSO. |
| Thaw Protocol |
| Partially immerse the vial in a 37°C water bath with gentle shaking until most of the medium is thawed. In a tissue culture hood, add 1 ml of pre-warmed culture medium into the vial and immediately transfer the contents of the vial to a centrifuge tube containing 5-10 ml of pre-warmed culture medium. Centrifuge the tube at room temperature for 5 minutes, aspirate the supernatant and suspend the cell pellet in 5-10 ml of pre-warmed culture medium. |
| Culture Protocol |
| RPMI8226-GFP-LUC is a cell line that grows in suspension. Culture the cells in RPMI-1640 medium containing 10% FBS using a humidified incubator set to 5% CO ₂ . When the culture has reached a density of 2 million cells per ml, add a 10-fold volume of fresh, pre-warmed culture medium to the cells. |
| Notices & Disclaimer |
| ProMab Biotechnologies products are intended for laboratory research purposes only, not for use in humans. This product is not for resale and may not be transferred to a third party without written consent from ProMab Biotechnologies, Inc. To purchase this product, you must accept the terms and conditions of ProMab Biotechnologies' Material Transfer Agreement (MTA). |

All products are for research use only

Discover more | ProMab.com



📍 2600 Hilltop Dr, Building B, Richmond, CA 94806
☎ 1.866.339.0871 | ✉ info@promab.com
📞 510.740.3625 | ✉ customerservice@promab.com






Figure 1. PM-RPMI8226-GFP-LUC cells and parental RPMI8226 cells were analyzed for GFP fluorescence by flow cytometry. Only the PM-RPMI8226-GFP-LUC cells were fluorescent.




Figure 2. PM-RPMI8226-GFP-LUC cells and parental RPMI8226 cells were lysed and incubated with luciferin to assess luciferase expression. Only PM-RPMI8226-GFP-LUC cells became luminescent.

All products are for research use only

Discover more | ProMab.com



2600 Hilltop Dr, Building B, Richmond, CA 94806
 1.866.339.0871 | info@promab.com
 510.740.3625 | customerservice@promab.com

